Application No.: 10/762,583 Docket No.: 1268-215

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

- 1. (Previously presented) A method for treatment of hard tissues present in a fluid-filled body cavity, wherein the fluid-filled body cavity is selected from salivary ducts and temporomandibular joints, the cavity having a diameter of 3 mm or less, the method comprising:
 - a) generating a laser beam using an Er:YAG laser device, said laser beam having a wavelength of about 2940nm; and
 - b) applying said laser beam to said hard tissue, or to a proximity of said hard tissue.
- 2. (Canceled)
- 3. (Original) A method according to Claim 1, wherein the hard tissue is fibrous scar tissue or calculi.
- 4. (Original) A method according to Claim 1, wherein the hard tissue is disintegrated to fragments having a size of less than 2 mm.
- 5. (Original) A method according to Claim 1, wherein the laser beam is provided through an endoscope, said endoscope also used for viewing the hard tissue.
- 6. (Original) A method according to Claim 5, wherein the endoscope is a Nahlieli type sialoendoscope.
- 7. (Original) A method according to Claim 1, wherein the parameters of the laser beam are 200-1000 millijoule/mm².
- 8. (Previously presented) A method according to claim 7, wherein the parameters of the laser beam are 300-700 millijoule/mm².

Application No.: 10/762,583 Docket No.: 1268-215

9. (Previously presented) A method according to claim 8, wherein the parameters of the laser beam are 500-700 millijoule/mm².

- 10. (Previously presented) A system for carrying out the method of Claim 1, said system comprising:
- (a) an endoscope for visualizing the interior of the cavity of said body cavity;
- (b) an Erbium laser device located in said endoscope, adapted to generate a laser beam in order to pulverize the hard tissue; and
- (c) an optic fiber for delivering the laser beam to the hard tissue or to the vicinity of the hard tissue, the length of the optic fiber being 10-20 cm.
- 11. (Original) The system according to Claim 10, wherein the endoscope is a Nahlieli type sialo-endoscope, and wherein said delivery of said laser beam is by a rigid, curved optical fiber.
- 12. (Previously presented) An endoscopic device comprising aperture adapted for connecting to an Er laser having an optic fiber for insertion into a body cavity having a diameter of 3mm or less.
- 13. (Previously presented) A device according to claim 12, having an optic fiber having a length of 10-20 cm.
- 14. (Previously presented). A device according to claim 13, wherein the optic fiber is flexible.
- 15. (Previously presented) A method for treatment of hard tissue present in any one of salivary ducts, temporomandibular joints and the like, comprising:
 - a) generating laser radiation using an Er:YAG laser device, said laser radiation having a wavelength of about 2940nm; and
 - b) irradiating said hard tissue with said laser radiation.
- 16. (Previously presented) The method of claim 15, wherein said hard tissue is in the form of salivary stones.

Application No.: 10/762,583 Docket No.: 1268-215

17. (Previously presented) The method according to claim 16, wherein said method is applied to any one of the sub-mandibular gland, sublingual gland, parotid duct, and parotid glands.

- 18. (Previously presented) The method according to claim 15, particularly for the treatment of scars, diseases and disorders in the temporomandibular joints.
- 19. (Previously presented) The method according to claim 15, wherein said laser radiation comprises an intensity of between about 200 and about 1000 millijoule/mm².
- 20. (Previously presented) The method according to claim 15, wherein said laser radiation is delivered in a pulsed manner.
- 21. (Previously presented) The method according to claim 15, wherein said laser radiation comprises an intensity of between about 300 and about 700 millijoule/mm².
- 22. (Previously presented) The method according to claim 15, wherein said laser radiation comprises an intensity of between about 500 and about 700 millijoule/mm².